



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\Delta = \begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix}, \quad \Delta' = \begin{vmatrix} a' & h' & g' \\ h' & b' & f' \\ g' & f' & c' \end{vmatrix}.$$

$$\Theta = Aa' + Bb' + Cc' + 2Ff' + 2Gg' + 2Hh',$$

$$\Theta' = A'a + B'b + C'c + 2F'f + 2G'g + 2H'h,$$

where A, B, C , etc., are minors of Δ , and A', B', C' , etc., are minors of Δ' .

$$\text{For } \frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 = 0 \text{ and } \frac{x^2}{a'^2} + \frac{y^2}{b'^2} - 1 = 0, \text{ we find } \Delta' = -\frac{1}{a'^2 b'^2},$$

$$\Theta = -\frac{1}{a^2 b'^2} - \frac{1}{a'^2 b^2} - \frac{1}{a^2 b^2}, \quad \Theta' = -\frac{1}{a'^2 b^2} - \frac{1}{a^2 b'^2} - \frac{1}{a'^2 b'^2}.$$

Substituting in (1), we have

$$\left(\frac{a'^2}{a^2} + \frac{b'^2}{b^2} + 1 \right)^2 = 4 \left(\frac{a'^2}{a^2} + \frac{b'^2}{b^2} + \frac{a'^2 b'^2}{a^2 b^2} \right)$$

$$\text{from which } \frac{a'}{a} \pm \frac{b'}{b} = \pm 1.$$

The only real solution for ellipses is

$$\frac{a'}{a} + \frac{b'}{b} = 1.$$

In problem 142,

$$a' = \frac{ab^2}{a^2 + b^2}, \quad b' = \frac{a^2 b}{a^2 + b^2}, \text{ from which } \frac{a'}{a} + \frac{b'}{b} = 1.$$

A very excellent demonstration was received from *G. B. M. ZERR*.

CALCULUS.

NOTE ON CENTER OF CURVATURE.

By *GEORGE R. DEAN, A. M.*, Professor of Mathematics, University of Missouri School of Mines and Metallurgy, Rolla, Mo.

The fact, that the point of intersection of two normals which approach each other is neither at infinity nor at the foot of the normals when they become coincident, is not plain to most students. The difficulty may be overcome, in some instances, in the following manner.

Let $(x_1, y_1), (x_2, y_2)$ be two points of the curve; $m_1; m_2$ the slope of the tangents at these points. The equations of the normals are then

$$x+m_1y=x_1+m_1y_1, \quad x+m_2y=x_2+m_2y_2.$$

It is proposed to find the coördinates of the intersection of these lines when (x_1, y_1) approaches and becomes coincident with (x_2, y_2) . Eliminating x between the equations, we have

$$y=y_1+\frac{1+m_2\left(\frac{y_1-y_2}{x_1-x_2}\right)}{\frac{m_1-m_2}{x_1-x_2}}.$$

Since $\text{Lim.}_{x_1=x_2}\left(\frac{y_1-y_2}{x_1-x_2}\right)=\frac{dy_1}{dx_1}$ and $\text{Lim.}_{x_1=x_2}\left(\frac{m_1-m_2}{x_1-x_2}\right)=\frac{d^2y_1}{dx_1^2}$, we have

$$y=y_1+\frac{1+\left(\frac{dy_1}{dx_1}\right)^2}{\frac{d^2y}{dx^2}}, \quad \text{and} \quad x=x_1-\frac{dy_1}{dx_1}\left\{\frac{1+\left(\frac{dy_1}{dx_1}\right)^2}{\frac{d^2y_1}{dx_1^2}}\right\}.$$

105. Proposed by CHARLES C. CROSS, Meridithville, Va.

From all points in a straight line passing through the center of a given circle tangents are drawn to the circle. If the bases and vertices of all the angles thus formed are made to coincide; required the equation of the curve passing through the tangent points.

Solution by J. W. YOUNG, Oliver Graduate Scholar in Mathematics, Cornell University, Ithica, N. Y.

Let the circle be $x^2+y^2=a^2$, and the given line $y=0$. Then the length of the tangent from any point $(x_1, 0)$ on the given line is $\sqrt{(x_1^2-a^2)}$.

Also the slope of the tangent may be calculated from the equation $mx_1 \pm a\sqrt{(1+m^2)}=0$, which is obtained by substituting the point $(x_1, 0)$ in the tangent equation $y=mx \pm a\sqrt{(1+m^2)}$. Solving for m , we have

$$m^2=\frac{a^2}{x_1^2-a^2}, \quad \text{or} \quad m=\pm\frac{a}{\sqrt{(x_1^2-a^2)}}.$$

To determine the required locus, use polar coördinates, with the common vertex of angles as pole and their common base as initial line; the coördinates (r, θ) of the tangent points are then given by the equations

$$\left. \begin{aligned} r &= \sqrt{(x_1^2-a^2)} \\ \tan\theta &= \frac{\pm a}{\sqrt{(x_1^2-a^2)}} \end{aligned} \right\}$$

Hence, the required locus is $r \tan\theta = \pm a$; in Cartesian coördinates this becomes $x^2(y^2-a^2)+y^4=0$.

Also solved by H. C. WHITAKER, and G. B. M. ZERR.